



Potoo

Potoos are a noisy, strange-looking bird, which can often be found in the canopy of the Amazon rainforest.

Appearance

- They have huge, gaping mouths.
- Potoos have large, sometimes yellow, eyes.
- Their feathers are grey, brown and black, and are perfect as camouflage against tree trunks.

Diet

 The potoo mostly eats bugs and insects, including grasshoppers, moths and beetles.



"Mrs. Moon" by julian londono is licensed under CC BY 2.0

- Potoos can be found in humid forests in Central and South America.
- They live in the high branches of trees, camouflaged among the tree bark.

Sloth

Sloths are nocturnal mammals and are known for being particularly slow-moving creatures.

Appearance

- Species of sloths are divided into two families: two-toed and three-toed sloths.
- They have thick fur, rounded heads, flat snouts and stumpy tails.
- · They commonly have algae growing on their fur.

Diet

- Some sloth species only eat leaves.
- Other types of sloth also eat fruit, insects and even small lizards.

"Crawling" by ndanger is licensed under CC BY 2.0

- Sloths are mostly found in Central and South American rainforests.
- They mainly live among the branches of trees.

Emerald Tree Boa

The emerald tree boa is a non-venomous snake, which is often found in the Amazon rainforest.

Appearance

- Emerald tree boas are bright green with a yellow underside.
- They have an irregular, zigzag pattern along their back.
- Adult emerald tree boas can grow to reach up to 2 metres.
- They have a pair of long, sharp teeth, pointed backwards.
- Their heads are relatively large and flat.

Diet

 They feed mostly on small mammals such as rats, bats and possums.



"Emerald Tree Boa" by Eric Kilby is licensed under CC BY 2.0

- Emerald tree boas live in the trees of the Amazon rainforest.
- They can also often be found near rivers or swamps but are not dependent on open water.

Red-Eyed Tree Frog

The red-eyed tree frog is just one of many species of frog that lives in the rainforest.

Appearance

- They are known for having huge, bright red eyes.
- The rest of their body is neon green with blue and yellow at the sides.
- They have specially adapted suction-cup toes which help them to cling on to tree branches and leaves.



"Red Eyed Tree Frog" by Douglas Tofoli is licensed under CC BY 2.0

Diet

The red-eyed tree frogs' prey includes moths, crickets, flies and grasshoppers.

Habitat

- They are commonly found in the tropical lowland rainforests or surrounding hills.
- They commonly inhabit areas close to rivers or ponds.

back

Scarlet Macaw

The scarlet macaw is one of at least 17 known species of macaws. Most are intelligent, social birds.

Appearance

- Despite their name, scarlet macaws are multicoloured: red, yellow and blue.
- They have large, powerful beaks.
- Their bodies are between 80cm to 90cm long.

Diet

 They mostly eat fruit and seeds or nuts, but can also eat insects.



"IMGP4701" by Kurayba is licensed under CC BY 2.0

- They thrive in the tropical, humid atmosphere of the rainforest.
- They are usually found in the emergent or canopy layers of the forest.

Jaguar

The jaguar is a member of the wild cat family that can be found in the rainforest regions.

Appearance

- They are famous for their beautiful, spotted coats.
- They can reach up to six feet in length, from their nose to the tip of their tail.
- They have powerful jaws and sharp teeth.



"Jaguar, Alert" by Eric Kilby is licensed under CC BY 2.0

Diet

 Jaguars are carnivores with an extensive list of possible meals including squirrels, deer and monkeys.

Habitat

- Jaguars are found in rainforests as well as grasslands and woodlands.
- They spend much of their time on the ground or the forest floor, using their soft padded paws to stalk quietly through the undergrowth.

back